

Hendra Virus

For the first time Australia, the lucky country, is faced with a deadly horse disease that is unique to our country. The fact that this disease can be caught by and kill humans as well, just increases the fear and worry that we all feel. It's no good saying that it's a rare event, because the devastation that it brings has a huge impact on equine owners, carers and industry.

Hendra virus was first isolated in 1994 from an outbreak of disease in a racing stable located in the northern Brisbane suburb of Hendra less than 10km from the city centre. The outbreak resulted in death of a horse trainer and 13 horses and left a stable hand seriously ill. A further seven horses with evidence of exposure to the virus were humanely destroyed to avoid possible further spread of the disease. Since Hendra virus was first isolated, significant progress has been made in understanding the virus, where it originates in nature, and how to detect infection and past exposure. Hendra virus and its close relation Nipah virus, both belong to a separate category of the Paramyxoviridae family, the Henipaviruses. In addition to a number of unique molecular characteristics, these viruses are distinguished from other members of the family by their ability to infect a broad range of species and fatally infect both animals and humans. Nipah virus emerged in pigs and humans in Malaysia. Since then, over 470 known human infections and over 240 deaths have been linked to outbreaks of Nipah in Malaysia, Singapore, Bangladesh and India. There have been no reported outbreaks in Australia.

So what does Hendra do – in very simple terms it attacks the cells so that they leak blood and fluids. Where it attacks in the body leads to the broad range of symptoms. Hendra cases have also exhibited anything from mild depression to colic – we can now say that there are no “classic” Hendra symptoms. All sick horses should be treated with caution. In most of the recorded infected cases, there has been strong presentation of clinical signs; however, occasional cases have demonstrated a much milder presentation of clinical signs.

What to look for!

Common Clinical Signs

- acute onset of illness
- increased body temperature
- increased heart rate
- discomfort/weight shifting between legs (both fore and hind limbs)
- depression
- rapid deterioration, usually with respiratory and/or neurological signs.

Some other clinical observations that have been noted include the following.

Respiratory signs include

- pulmonary oedema and congestion
- respiratory distress—increased respiratory rates
- terminal nasal discharge—can be initially clear progressing to stable white froth and/or stable blood stained froth

Neurological signs that include:

- ‘wobbly gait’ progressing to ataxia
- altered consciousness—apparent loss of vision in one or both eyes, aimless walking in a dazed state
- head tilting, circling
- muscle twitching— spasms have been seen in acutely ill and recovered horses
- urinary incontinence
- lying down with inability to rise.

Other possible signs

- previous unexplained horse deaths
- facial oedema (swelling)
- facial paralysis and/or a locked jaw
- spasms of the jaw, involuntary chomping
- muscle trembling
- altered gait, high stepping
- anorexia
- congestion of oral mucous membranes
- a high case fatality rate within 48 hours where there are multiple cases
- colic-like symptoms in some cases (generally quiet abdominal sounds on auscultation of the abdomen in pre-terminal cases)
- straining with difficulty passing manure
- stranguria (difficult urination) or dribbling urine in both males/females, and hot hooves
- bad breath/halitosis
- delayed blood clotting times.

**If you see these symptoms call your veterinarian or
 DPI&F 13 25 23 or Emergency Animal Disease
 1800675888 Shower and change clothes if you have
 handled a sick horse - Stay away from other horses**

**If you have a horse that is suspected to
 have Hendra, do not move any other horses
 off the property until given the all clear.**

Horse with Hendra Virus - Biosecurity is Essential



Biosecurity must be carried out!

Hendra is quite hard to catch but it is one of the deadliest viruses known. At this stage there are no effective treatments. Please read our fact sheets on [Biosecurity and Hendra Virus – Reducing the Risk](#)

The following points when combined with the symptoms already listed could also support suspicion of HeV:

- Where there are multiple cases, a high case fatality rate occurs within 48 hours.
- Some cases have initially been reported as colic or even depression – there are no “classic” Hendra symptoms.
- Bats in the area, though a lack of bat sightings does not rule out HeV.
- The animals live in or near paddocks that attract bats. These include trees with soft fruits, figs, stone fruits, or flowering trees with high nectar or pollen

For more information on property design and safer trees to plant read Qld Horse Council’s fact Sheets [Hendra – Property Design](#) and [Bats & Trees](#)

Hendra virus is carried by flying foxes. Under unknown but rare circumstances, the virus spills over from these bats to equines (horses/donkeys etc), killing over 70% of the horses it infects. Under even rarer circumstances, the virus spreads to humans who have had very close contact with Hendra infected horses. While there is strong evidence to support this mode of transmission (bat-to-horse-to-human) there is no evidence of bat-to-human or human-to-human.

Though there have been significant gains in knowledge about Hendra, a great deal remains to be learnt. The incubation period (time from exposure to the appearance of the first clinical signs of infection) of Hendra virus in horses is five to 16 days. Fatally infected horses die on average two days after the first sign of

infection. While approximately 25% of horses are thought to survive acute infection, the current national policy requires these horses to be euthanased. The incubation period in humans is believed to be five to 21 days, and the current human case fatality rate is more than 50%.

Human Hendra Virus Infections

Mackay, August 1994. Not recognised until the death of the farmer who had assisted his veterinarian wife with an autopsy of two horses that died of unknown causes. The farmer appeared fully recovered after hospitalisation but relapsed and died 13 months later. Retrospective testing showed the horses were infected with Hendra.

Hendra, September 1994. Resulted in the death of a horse trainer, 13 of his horses, and caused severe febrile illness in a stable-hand. The trainer had very close physical contact sick horses, as did the stable hand.

Gordonvale, October 2004. A veterinarian tested positive for Hendra virus after performing a post mortem on a horse that died suddenly. While no samples were available from the horse for testing, the clinical and post mortem signs were consistent with Hendra virus infection.

Redlands, June 2008. A veterinarian and veterinary nurse at the Redlands Veterinary Clinic were infected after close direct contact with infected horses. The Veterinarian tested positive for Hendra virus and died in August after spending several weeks in intensive care. The veterinary nurse also spent several weeks in intensive care but was later discharged.

Cawarral, August 2009. A Queensland veterinarian died in September after being exposed to Hendra. Infection has been confirmed in two horses that died and another had to be destroyed after returning a positive test. The incident was discovered after a property manager and a local veterinarian alerted Biosecurity Queensland with the death of a horse suffering from respiratory distress.

How is the virus excreted?


The detection of viral genetic material in the blood, nasal secretions and a wide range of body tissues of infected horses indicates that by the time a horse shows clinical signs of infection the virus is widespread throughout the body. Most virus is shed from these horses when they are sickest, suggesting that this is the most likely time for transmission to humans. However, studies have also shown that a horse can potentially excrete the virus through nasal/naso-pharyngeal secretions at least two days prior to the appearance of clinical signs.

Transmission

The exact mechanics of Hendra virus transmission from flying foxes to horses is not known. However transmission is thought to be through the ingestion of grass, feed or partially eaten fruit contaminated with bat urine, saliva or other bodily fluids.

Horse-to-horse transmission of the virus can happen, as a proportion of incidents involved infection of both the index case and companion horses. When a Hendra incident occurs other animals are tested on the Infected Properties, including all companion animals such as cats and dogs. To date there has only been one case of a dog showing antibodies but no signs of infection or disease. This mirrors work done in laboratories which showed that dogs were not susceptible but did develop antibodies.

The migratory and feeding areas of flying foxes



The Qld Horse Council stresses the need to be aware that Hendra virus can occur wherever there are flying foxes and horses, and because of the large area that flying foxes travel over, can occur across a large proportion of the state. .

How to predict your risks? You Can't!

- Hendra events have been rare but are increasing in frequency. Is that awareness and consequently more testing and detection? Have there previously been undetected cases?
- Proximity to Flying Fox roost has not been a significant factor in the location of infected properties. Flying foxes travel large distances to feed, including across your property.
- All Hendra events in the same calendar year have been large distances apart. They are not the same Flying Foxes causing the spillover.
- Do not rely on "seasonal" – the stressors that lead to bats excreting Hendra are not understood, and even if they were we have no way of controlling them. (drought/pregnancy/Flood?)
- All Hendra events have been of a random nature and the properties that have been affected have been different. (Topography/type of trees)

Table of all **KNOWN** incidents of Hendra Virus

Location	Month	Year	Horse Deaths	Human Cases	Human Deaths
Mackay	August	1994	2	1	1
Hendra	September	1994	20	2	1
Cairns	January	1999	1	0	0
Cairns	November	2004	1	1	0
Townsville	December	2004	1	0	0
Peachester	June	2006	1	0	0
Murwillumbah	October	2006	1	0	0
Peachester	June	2007	1	0	0
Clifton Beach	July	2007	1	0	0
Redlands	June-July	2008	5	2	1
Proserpine	July	2008	3	0	0
Cawarral	August	2009	4	1	1
Bowen	September	2009	2	0	0
Tewantin	May	2010	1	0	0
Boonah	June/July	2011	1 dog 3	0	0
Beaudesert	June	2011	1	0	0
Wollongbar	June	2011	1	0	0
Parkridge	June	2011	1	0	0
Logan	June	2011	1	0	0
Macksville	July	2011	1	0	0
Kuranda	July	2011	1	0	0
Boondall	July	2011	1	0	0
Hervey Bay	July	2011	1	0	0
Lismore	July	2011	1	0	0
Chinchilla	July	2011	1	0	0
Mullumbimby	July	2011	1	0	0
Ballina	August	2011	1	0	0
South Ballina	August	2011	2	0	0
Mullumbimby	August	2011	1	0	0
Gold Coast	August	2011	1	0	0
Ballina	August	2011	1	0	0
Beachmere	September	2011	2	0	0
Townsville	January	2012	1	0	0
Rockhampton	May	2012	1	0	0
Ingham	May	2012	1	0	0
Mackay	June	2012	1	0	0
Rockhampton	July	2012	3		
Cairns	July	2012	1		
Total			74	7	4

There's no BS in Biosecurity!

- Insist your vet use proper biosecurity and infection control practices.
- Every property must have a PPE Kit – contents are at any hardware store
- Reduce exposure to body fluids
- Protect sensitive areas (mucus membranes, eyes, wounds etc)
- Learn what to do if you suspect you are exposed.